

Space Weather Training Course



e-swan

SWEC
Space Weather Education Centre


SPACE WEATHER
Introduction




I want to present Dries.




On May 15, 2024, Dries is doing an HF test with his Canadian colleagues.
But the signal doesn’t pass through

What happened?



https://svs.gsfc.nasa.gov/5240


HF com - radio waves are being reflected in the ionosphere
GNSS signals pass through the ionosphere
Depends on f and fp :
f<fp reflection
f>fp transmission



https://svs.gsfc.nasa.gov/5240


The ionosphere consists of a gas which is, under the influence of the sun, partially ionised.

It is due to the ionisation that radio waves are being reflected or pass through, being absorbed, delayed.
The electrons decide which waves are reflected, pass through, how much they are absorbed and delayed.




https://svs.gsfc.nasa.gov/5240

The sun is the main factor/has a key role on the creation of the ionosphere.
Its energy can ionise the atmosphere.



https://svs.gsfc.nasa.gov/5240

We have the regular, foreseen solar weather.
Light - wind of plasma - energetic particles




https://svs.gsfc.nasa.gov/5240

This energy can evolve into storms busting down on Earth

Solar flare – solar wind disturbances – particle storms


Space Weather
The Sun's energy impacting earth's atmosphere and magnetosphere



As we go out into space, the atmosphere becomes very thin, until by the time we are in space, it has almost vanished. Almost, but not quite. Even in space there are some atoms which are often moving very quickly. Many forms of energy also move through **space** and it is the **interaction of energy and atoms that produces what we refer to as space weather**. In particular, space weather is the changes that occur in the space environment.

The **sun** is the source of 'normal' terrestrial weather. It is also the **primary (but not the only) source of space weather**. Most aspects of space weather affect us to some extent. The more our society becomes dependent on technology and the more we utilize space, the more we are affected by space weather. Some aspects of space weather are benevolent, and allow activities not otherwise possible such as long range radio communications. Some aspects are benign but fascinating such as the Aurora, and some are malevolent. **Like terrestrial weather, it depends on the situation and the event.**




The sun is a gigantic ball of energy: magnetic energy, heat, moving plasma, …

This energy is kept inside the Sun but also on its surface and in its atmosphere in magnetic structures like sunspots and magnetic loops, filaments or prominences ready to be released.

This energy is expelled, leaves the Sun to outer space and is carried away by electromagnetic waves, plasma and energetic particles.

Note: the solar plasma is hot. The plasma particles bump on each other. These collisions changes their kinetic energy. This change is emitted in the form of thermal radiation, light photons. Once these photons are at the solar surface, they can escape and move freely.
Thermal radiation is electromagnetic radiation generated by the thermal motion of particles in matter. You have thermal motion as soon as the temperature is above absolute zero.




100 kEV
Energetic particles detach from the plasma and move on their own

Energetic subatomic particles are individual constituents of atoms, such as electrons, protons, and neutrons, that are moving at high speeds or have high kinetic energy. Plasma, in contrast, is a distinct state of matter consisting of a superheated, ionized gas where these subatomic particles (now called ions and free electrons) are separated and move freely, rather than being bound within neutral atoms. Therefore, energetic subatomic particles are the fundamental building blocks, while plasma is the collective, highly energized, and ionized state formed by these particles.

3 SPACE WEATHER PHENOMENA

The sun's energy reaches the earth in 3 forms: light, moving gas and particle precipitation. This energy interacts with the magnetosphere and the atmosphere of the earth. This is space weather.

How and where the interaction occurs depends on the type of energy.



light= electromagnetic waves, energy transmitted in the form of photons/ EM waves

Wind= moving gas, in this case moving plasma

EPP = particle has high energy and precipitates along the geomagnetic field.


<100 kEV : plasma

Particle precipitation - energy transmitted in the form of fast-moving atomic or sub-atomic particles —> drizzle

Energy expressed in eV= 1.602 10^-19 J


https://lasp.colorado.edu/home/mag/research/energetic-particle-precipitation/

Focus on solar energetic particles




Magnetism drives space weather and the sun’s dynamics.

In the EUV it looks like this!
The sun doesn’t only rotate but it is also tangled with loops which move, shake back and forth, restructure and do all sorts of electric and magnetic magic.
It is a turbulent plasma ball where magnetic fields are moving, shake, connect with other other, ….



Sunspots are bundles of magnetic fields.
This is the photosphere of the Sun, in the visible light.





The more sunspots, the higher the chance for magnetic reconnection.

0.3 T – solar sunspot
5mT – strength of a typical refrigerator magnet
31.869 μT (3.1×10^{-5} T) – strength of Earth's magnetic field at 0° latitude (North/South), 0° longitude (west/east)
1 to 5 nT – IMF at L1

SOLAR WEATHER & STORMS

Energy can be released on a shorter time scale. A solar feature like a sunspot, an active region, coronal hole, filament etc. lies at the base of a solar storm in which energy is released. The release of energy might be in an abrupt, impulsive and brutal way (flare, Coronal Mass Ejection or CME, proton storm) or in a non-eruptive manner (Coronal Hole - CH).





Change in energy output on the scale of minutes, hours, days.

Remote sensing (seeing) - in situ (taste and touch the ambient space)

Space weather is the change of energy that occur in the space environment.

A Flare is a sudden strong increase of the solar e.m. radiation. The light flash is localised on the solar surface.
SDO/AIA

A Coronal Mass Ejection is a plasma cloud that is ejected into space. You consider it as a cloud and not as a bunch of individual particles. It is superimposed on the background solar wind. You can see a CME as a complex magnetic bag with different magnetic layers with plasma in it that travels as a tsunami through space. It can go faster/as fast as/slower than the background solar wind. When it is faster, you will see a shock in front of the cloud. This is exactly the same as the shock you see in front of a speed boat.
A CME is visible as a white cloud in corona graphic images like the one on the slide. A coronagraph is a telescope that creates an artificial eclipse and makes pictures in the visible light of the region around the sun.
SOHO/LASCO C2 (red) and LASCO C3 (blue)


A coronal hole is a structure in the solar corona that you see as a black area in the EUV. It looks black because there is less plasma present that radiates in the EUV. The magnetic field lines are open, i.e. fan out into space. There are no magnetic loops above a coronal hole. The solar wind emanating from a CH is faster compared to the usual solar wind.
SDO/AIA

Particle shower
A particle storm is a bunch of electrically charged particles that are accelerated in the solar atmosphere to very high velocities by a large-scale magnetic eruption often causing a CME and/or solar flare. They follow the IMF
They may impact telescopes. They are seen as white stripes and dots: this are particles that fall into the lens and blind the pixel(s). During that particular moment, the telescope can't see anymore through the impacted pixels. You can say that the dots and stripes represent a sort of in situ measurement.

In situ means that you measure a parameter local. Remote sensing means that you look at something from a distance.

Near Earth, the IMF still controls the solar wind and its movement. If we would go much much further, the CME magnetic bag with solar plasma would be almost empty (all the solar material is spread

over an immense volume) and the magnetic bag would have evaporated. But, this doesn't matter for us. We are at 1AU and at 1AU the IMF and solar plasma make space weather in a normal way, in an extreme way.



Change in energy output on the scale of minutes, hours, days.

Remote sensing (seeing) - in situ (taste and touch the ambient space)

Space weather is the change of energy that occur in the space environment.

A Flare is a sudden strong increase of the solar e.m. radiation. The light flash is localised on the solar surface.
SDO/AIA

A Coronal Mass Ejection is a plasma cloud that is ejected into space. You consider it as a cloud and not as a bunch of individual particles. It is superimposed on the background solar wind. You can see a CME as a complex magnetic bag with different magnetic layers with plasma in it that travels as a tsunami through space. It can go faster/as fast as/slower than the background solar wind. When it is faster, you will see a shock in front of the cloud. This is exactly the same as the shock you see in front of a speed boat.
A CME is visible as a white cloud in corona graphic images like the one on the slide. A coronagraph is a telescope that creates an artificial eclipse and makes pictures in the visible light of the region around the sun.
SOHO/LASCO C2 (red) and LASCO C3 (blue)

A coronal hole is a structure in the solar corona that you see as a black area in the EUV. It looks black because there is less plasma present that radiates in the EUV. The magnetic field lines are open, i.e. fan out into space. There are no magnetic loops above a coronal hole. The solar wind emanating from a CH is faster compared to the usual solar wind.
SDO/AIA

Particle shower
A particle storm is a bunch of electrically charged particles that are accelerated in the solar atmosphere to very high velocities by a large-scale magnetic eruption often causing a CME and/or solar flare. They follow the IMF
They may impact telescopes. They are seen as white stripes and dots: this are particles that fall into the lens and blind the pixel(s). During that particular moment, the telescope can't see anymore through the impacted pixels. You can say that the dots and stripes represent a sort of in situ measurement.

In situ means that you measure a parameter local. Remote sensing means that you look at something from a distance.


Near Earth, the IMF still controls the solar wind and its movement. If we would go much much further, the CME magnetic bag with solar plasma would be almost empty (all the solar material is spread

over an immense volume) and the magnetic bag would have evaporated. But, this doesn't matter for us. We are at 1AU and at 1AU the IMF and solar plasma make space weather in a normal way, in an extreme way.



Transit time

On earth
Storm scale – strength
duration
Impact
Area of impact



Light storms

During a flare, an area in the solar corona lights up. This is a movie from the EUV imager AIA onboard of SDO.

AU TRANSIT TIME

The energy released during a solar storm moves through space, each with its own typical speed: speed of light, ….



AU TRANSIT TIME

The energy released during a solar storm moves through space, each with its own typical speed: speed of light, …



8 MIN



STORM SCALE

https://www.spaceweather.gov/products/goes-x-ray-flux




The scale of a flare is defined by it’s X-ray flux. The X-ray flux is measured by the geostationary satellite GOES.

STORM SCALE



A,B,C,M,X




https://www.spaceweather.gov/products/goes-x-ray-flux

The scale of a flare is defined by it’s X-ray flux. The X-ray flux is measured by the geostationary satellite GOES.

DURATION

MINs to HOUR



IMPACT





The icon represent the Earth. White is the day-side, black is the night side.
When you ‘see’/detect a flare, you can be impacted.

Electromagnetic radiation

- Photons / electromagnetic waves
- Speed of light




Photo-ionisation — green


Ionizing radiation is a type of energy released by atoms that travels in the form of electromagnetic waves (gamma or X-rays) or particles (neutrons, beta or alpha). The spontaneous disintegration of atoms is called radioactivity, and the excess energy emitted is a form of ionizing radiation.

Ionizing radiation (or ionising radiation), including nuclear radiation, consists of subatomic particles or electromagnetic waves that have sufficient energy to ionize atoms or molecules by detaching electrons from them.[1] Some particles can travel up to 99% of the speed of light, and the electromagnetic waves are on the high-energy portion of the electromagnetic spectrum.

Gamma rays, X-rays, and the higher energy ultraviolet part of the electromagnetic spectrum are ionizing radiation, whereas the lower energy ultraviolet, visible light, nearly all types of laser light, infrared, microwaves, and radio waves are non-ionizing radiation. The boundary between ionizing and non-ionizing radiation in the ultraviolet area cannot be sharply defined, as different molecules and atoms ionize at different energies. The energy of ionizing radiation starts between 10 electronvolts (eV) and 33 eV.

IMPACT










The icon represent the Earth + atmosphere with the ionosphere
Ionisation of atoms and molecules due to photons -

The light smashes atomic particles into pieces. Those pieces are electrically charged.

TRANSPARENCY OF EARTH ATMOSPHERE

Our atmosphere protects us against harmful solar light. This ionising solar light can penetrate only until a certain height.




AREA OF IMPACT



AREA OF IMPACT












The icon represent the Earth. White is the day-side, black is the night side.
When you ‘see’/detect a flare, you can be impacted.





Change in energy output on the scale of minutes, hours, days.

Remote sensing (seeing) - in situ (taste and touch the ambient space)

Space weather is the change of energy that occur in the space environment.

A Flare is a sudden strong increase of the solar e.m. radiation. The light flash is localised on the solar surface.
SDO/AIA

A Coronal Mass Ejection is a plasma cloud that is ejected into space. You consider it as a cloud and not as a bunch of individual particles. It is superimposed on the background solar wind. You can see a CME as a complex magnetic bag with different magnetic layers with plasma in it that travels as a tsunami through space. It can go faster/as fast as/slower than the background solar wind. When it is faster, you will see a shock in front of the cloud. This is exactly the same as the shock you see in front of a speed boat.
A CME is visible as a white cloud in corona graphic images like the one on the slide. A coronagraph is a telescope that creates an artificial eclipse and makes pictures in the visible light of the region around the sun.
SOHO/LASCO C2 (red) and LASCO C3 (blue)


A coronal hole is a structure in the solar corona that you see as a black area in the EUV. It looks black because there is less plasma present that radiates in the EUV. The magnetic field lines are open, i.e. fan out into space. There are no magnetic loops above a coronal hole. The solar wind emanating from a CH is faster compared to the usual solar wind.
SDO/AIA

Particle shower
A particle storm is a bunch of electrically charged particles that are accelerated in the solar atmosphere to very high velocities by a large-scale magnetic eruption often causing a CME and/or solar flare. They follow the IMF
They may impact telescopes. They are seen as white stripes and dots: this are particles that fall into the lens and blind the pixel(s). During that particular moment, the telescope can't see anymore through the impacted pixels. You can say that the dots and stripes represent a sort of in situ measurement.

In situ means that you measure a parameter local. Remote sensing means that you look at something from a distance.

Near Earth, the IMF still controls the solar wind and its movement. If we would go much much further, the CME magnetic bag with solar plasma would be almost empty (all the solar material is spread


over an immense volume) and the magnetic bag would have evaporated. But, this doesn't matter for us. We are at 1AU and at 1AU the IMF and solar plasma make space weather in a normal way, in an extreme way.



A CME is visible as a white cloud in corona graphic images like the one on the slide. A coronagraph is a telescope that creates an artificial eclipse and makes pictures in the visible light of the region around the sun.
SOHO/LASCO C2


AU TRANSIT TIME


The energy released during a solar storm moves through space, each with its own typical speed: speed of light, order of a few 100 km/s,….



AU TRANSIT TIME

The energy released during a solar storm moves through space, each with its own typical speed: speed of light, order of a few 100 km/s, relativistic speeds.






A CME that hits the Earths magnetosphere.
Precipitating electrons coming from the tail of the magnetosphere gyrate along the earths magnetic field and drop into the atmosphere in the auroral oval.

These electrons have no solar origin, they are present in the plasmasphere of the Earth.


STORM SCALE

https://kp.gfz.de/en/



The Kp index is an index that quantifies the disturbance of the magnetic field of Earth. It ranges between 0 and 9, with 0 no disturbance and 9 an extreme disturbance.

STORM SCALE




The Kp index is an index that quantifies the disturbance of the magnetic field of Earth. It ranges between 0 and 9, with 0 no disturbance and 9 an extreme disturbance.


measure for the energy input from the solar wind to Earth and it is used by space weather services in near real-time. Kp goes back to 1932 and is an important parameter to investigate long-term climate change in the upper atmosphere, in geospace, in the solar wind and on the sun. Kp is based on the data of 13 geomagnetic observatories around the globe.

https://kp.gfz.de/en/

STORM SCALE









The Kp index is an index that quantifies the disturbance of the magnetic field of Earth. It ranges between 0 and 9, with 0 no disturbance and 9 an extreme disturbance.


DURATION








Day or more



The icon represent the Earth. White is the day-side, black is the night side.
When you ‘see’/detect a flare, you can be impacted.




- Ampères law : circulation of magnetic field = enclosed electric current —> sheet current

A magnetic field is imbedded in the solar wind. This magnetic field can interact with the magnetic field of the earth at the boundaries of the earth magnetosphere. This interaction is called reconnection. It happens when 2 magnetic regions are confronted with each other.
The magnetic field carried by the solar wind can couple with the magnetic field of Earth. This coupling is stronger when the solar wind magnetic field is opposite to the magnetic field of Earth.

The blue magnetic field lines are imbedded in the solar wind.
The red magnetic field lines represent the earth magnetosphere.
The blue and the red magnetic region have to face each other. Opposite magnetic field lines can connect easily and ‘open’. This causes geomagnetic storms. Magnetic field lines in the same direction interact less.

0.3 T – solar sunspot
5mT – strength of a typical refrigerator magnet
31.869 μT (3.1×10^{-5} T) – strength of Earth's magnetic field at 0° latitude (North/South), 0° longitude (west/east)
1 to 5 nT – IMF at L1



currents

- Ampères law : circulation of magnetic field = enclosed electric current —> sheet current
- Magnetopause current, neutral sheet current, magnetotail current
- can propagate to the ionosphere (field aligned currents)

Ionosphere: a region with free electrons due to ionisation - conductive system

Magnetosphere-ionosphere coupling
The gas-system of the ionosphere is connected with magnetic system of Earth. Both systems intersect and interact with each other.
(Magnetopause current couples to pedderson and hall currents in low ionosphere (E and D-layer))
—> due to the coupling with a magnetic dipolar structure, the ionospheric behaviour is different at low, middle and high latitudes.

Field aligned currents along the magnetic field lines, connect the magnetospheric currents with ionospheric currents. The Pedersen and Hall currents are two main currents in the lower ionosphere. They are localized in the ionospheric D and E regions.

Charged particles can move rather freely along field lines and therefore are good electric conductors. Electric currents flow along the field lines and connect magnetosphere and ionosphere. Therefore every electric feature in the magnetosphere has an “image” in the ionosphere, and conversely. We identify the following regions:

Magnetopause ↔ footpoints of the cusps
Tail lobes ↔ polar caps
Plasma sheet ↔ auroral oval
Plasmasphere ↔ ionosphere at low latitude


The ionosphere is that part of the upper atmosphere where free electrons occur in sufficient density to have an appreciable influence on the propagation of radio frequency electromagnetic waves. This ionization depends primarily on the Sun and its activity. ionospheric structures and peak densities in the ionosphere vary greatly with time (sunspot cycle, seasonally, and diurnally), with geographical location (polar, auroral zones, mid-latitudes, and equatorial regions), and with certain solar-related ionospheric disturbances.


The major part of the ionization is produced by solar X-ray and ultraviolet radiation and by corpuscular radiation from the Sun. The most noticeable effect is seen as the Earth rotates with respect to the Sun; ionization increases in the sunlit atmosphere and decreases on the shadowed side. Although the Sun is the largest contributor toward the ionization, cosmic rays make a small contribution. Any atmospheric disturbance affects the distribution of the ionization.

The ionosphere is a **dynamic system controlled by** many parameters including **acoustic motions of the atmosphere, electromagnetic emissions, and variations in the geomagnetic field**. Because of its extreme sensitivity to atmospheric changes, the ionosphere is a very sensitive monitor of atmospheric events.

The most accurate way of measuring the ionosphere is with a ground-based ionosonde, which records data as ionograms.

AREA OF IMPACT






Latitude

The magnetic field carried by the solar wind can reconnect on the day side with the magnetic field of Earth.
On the night side, magnetic reconnection between opposite magnetic field of the magnetosphere of Earth. (No solar magnetic field involved.)

The icon of the dipole represents the Earth dipolar field.

The result is that the Earth’s magnetic field is disturbed and most strong in the polar regions.


IMPACT





Plasmasphere - ionosphere
Moving plasma —> VTEC - irregularities





Change in energy output on the scale of minutes, hours, days.

Remote sensing (seeing) - in situ (taste and touch the ambient space)

Space weather is the change of energy that occur in the space environment.

A Flare is a sudden strong increase of the solar e.m. radiation. The light flash is localised on the solar surface.
SDO/AIA

A Coronal Mass Ejection is a plasma cloud that is ejected into space. You consider it as a cloud and not as a bunch of individual particles. It is superimposed on the background solar wind. You can see a CME as a complex magnetic bag with different magnetic layers with plasma in it that travels as a tsunami through space. It can go faster/as fast as/slower than the background solar wind. When it is faster, you will see a shock in front of the cloud. This is exactly the same as the shock you see in front of a speed boat.
A CME is visible as a white cloud in corona graphic images like the one on the slide. A coronagraph is a telescope that creates an artificial eclipse and makes pictures in the visible light of the region around the sun.
SOHO/LASCO C2 (red) and LASCO C3 (blue)


A coronal hole is a structure in the solar corona that you see as a black area in the EUV. It looks black because there is less plasma present that radiates in the EUV. The magnetic field lines are open, i.e. fan out into space. There are no magnetic loops above a coronal hole. The solar wind emanating from a CH is faster compared to the usual solar wind.
SDO/AIA

Particle shower
A particle storm is a bunch of electrically charged particles that are accelerated in the solar atmosphere to very high velocities by a large-scale magnetic eruption often causing a CME and/or solar flare. They follow the IMF
They may impact telescopes. They are seen as white stripes and dots: this are particles that fall into the lens and blind the pixel(s). During that particular moment, the telescope can't see anymore through the impacted pixels. You can say that the dots and stripes represent a sort of in situ measurement.

In situ means that you measure a parameter local. Remote sensing means that you look at something from a distance.

Near Earth, the IMF still controls the solar wind and its movement. If we would go much much further, the CME magnetic bag with solar plasma would be almost empty (all the solar material is spread

over an immense volume) and the magnetic bag would have evaporated. But, this doesn't matter for us. We are at 1AU and at 1AU the IMF and solar plasma make space weather in a normal way, in an extreme way.




Particle showers

You see energetic particles that impact the telescope LASCO/C3 onboard of SOHO. They are seen as white stripes and dots: these are particles that fall into the lens, hit a pixel or more pixels. The impacted pixel in blinded. The dots and stripes represent an in situ measurement.

(The image in the middle of the occulter is an EUV image from the instrument AIA onboard of SDO.)


AU TRANSIT TIME

The energy released during a solar storm moves through space, each with its own typical speed: speed of light, order of a few 100 km/s, relativistic speeds.




AU TRANSIT TIME

The energy released during a solar storm moves through space, each with its own typical speed: speed of light, order of a few 100 km/s, relativistic speeds.




STORM SCALE


https://www.spaceweather.gov/products/goes-proton-flux




The GOES satellite measures the proton flux.
Storm: 10 pfu (proton flux units) for >10MeV
Major storm: 100 pfu for >100MeV




https://www.spaceweather.gov/products/goes-proton-flux


GOES Proton Flux (5-minute data)
6 Hour
1 Day
3 Day
7 Day
10³
10²
10¹
10⁰
10⁻¹
10⁻²
Particles - cm⁻² · s⁻¹ · sr⁻¹
CMPC 10 MeV Warning Threshold
00:00 Nov 11
12:00
00:00 Nov 12
12:00
00:00 Nov 13
12:00
00:00 Nov 14
12:00
00:00 Nov 15
12:00
00:00 Nov 16
12:00
00:00 Nov 17
12:00
00:00 Nov 18
Universal Time
GOES-18 ≥ 10 MeV
GOES-18 ≥ 50 MeV
GOES-18 ≥ 100 MeV
GOES-18 ≥ 500 MeV
Updated 2025-11-17 0555 UTC
Space Weather Prediction Center

STORM SCALE






GOES

The GOES satellite measures the proton flux.
Storm: 10 pfu (proton flux units) for >10MeV
Major storm: 100 pfu for >100MeV

DURATION






MINs to HOUR

DAY+

HOURS to DAYS

It takes in the order of an hour to reach Earth but the particle shower on Earth can last for days

IMPACT







Photo-ionisation — green


Ionizing radiation is a type of energy released by atoms that travels in the form of electromagnetic waves (gamma or X-rays) or particles (neutrons, beta or alpha). The spontaneous disintegration of atoms is called radioactivity, and the excess energy emitted is a form of ionizing radiation.

Ionizing radiation (or ionising radiation), including nuclear radiation, consists of subatomic particles or electromagnetic waves that have sufficient energy to ionize atoms or molecules by detaching electrons from them.[1] Some particles can travel up to 99% of the speed of light, and the electromagnetic waves are on the high-energy portion of the electromagnetic spectrum.

Gamma rays, X-rays, and the higher energy ultraviolet part of the electromagnetic spectrum are ionizing radiation, whereas the lower energy ultraviolet, visible light, nearly all types of laser light, infrared, microwaves, and radio waves are non-ionizing radiation. The boundary between ionizing and non-ionizing radiation in the ultraviolet area cannot be sharply defined, as different molecules and atoms ionize at different energies. The energy of ionizing radiation starts between 10 electronvolts (eV) and 33 eV.

IMPACT


Ionisation
Particle content
Ionisation
Particle Radiation




Latitude


The magnetic field carried by the solar wind can reconnect on the day side with the magnetic field of Earth.
On the night side, magnetic reconnection between opposite magnetic field of the magnetosphere of Earth. (No solar magnetic field involved.)

The icon of the dipole represents the Earth dipolar field.


The result is that the Earth’s magnetic field is disturbed and most strong in the polar regions.



The particles grab the open magnetic field lines and enter in this way the magnetosphere and finally the atmosphere of earth.



The higher the energy, the deeper they can penetrate into the Earths atmosphere.


8 min
X-ray: A,B,C,M,X
Min to Hour
Ionisation
Illuminated area
Days
Kp: 0 - 9
Day+
Ionisation
Particle content
Auroral Oval & more
Latitude
Hour
Proton flux:
(Major) Storm
Hours to Days
Ionisation
Particle radiation
Polar Caps
High Altitude

X-ray Flux Passport



Type

X-ray output (0.1-0.8 nm and 0.05-0.4 nm) of the solar spectrum versus time.

Use

To quantify solar flares happening on the side of the Sun facing Earth.

Units

Integrated over 1 min

$W \cdot m^{-2}$

Data location

https://www.swpc.noaa.gov/products/goes-x-ray-flux

Plot

X-axis: time in UTC
Y-axis: X-ray flux in W/m^2, i.e. how much energy (Joule) passes in 1 sec (W=Watt=J/s) through 1 square meter at geostationary height.

Classes

The long wavelength (0.1-0.8 nm, red and orange) is the reference for identification of the class. The red is measured by GOES 18, orange by GOES 16.
The flare classes are indicated on the right of the graphs. When the peak of the red/orange curve is in blue/green/yellow/orange/red, it is a A/B/C/M/X-class flare.

Senso

GOES-16
GOES-18


Senso

Geostation

Caden

Integrated

Kp-index Passport



What
Kp-index (NOAA) represents the planetary geomagnetic condition or the state of the magnetic fi
globe.

Use
To quantify the disturbance of the magnetic field of Earth. The scale ranges from 0 to 9, with 0 in
of the Earth magnetic field .

Plot
X-axis: data and time in UTC, time stamp every 3 hours. Ranges from 3 days in the left (left) to r
Y-axis:
coloured bars of 3 hours wide, height= Kp-index (NOAA)
Height of a bar represents a number: 0, 0+, 1-,1,1+, 2-,2,2+, … 8-,8,8+, 9-,9.

Estimated
by NOAA, US

Data location
https://www.swpc.noaa.gov/products/planetary-k-index

Definitive

The values become definitive after calculation and confirmation by the German Research Centre for Geosciences, Potsdam.

Magnetometers


The calculation is based upon the measurements done by magnetometers.

Observatories

Thirteen (13) Geomagnetic Observatories between 44 degrees and 60 degrees northern or southern geomagnetic latitude.

SWEC Space Weather Education Center

K Belgium Passport



What

K Belgium represents the geomagnetic conditions or the state of the magnetic field of Earth above Belgium. It is an index for Belgium.

Use

To quantify the disturbance of the magnetic field of Earth. The scale ranges from 0 to 9, with 0 indicating no disturbance and 9 the highest disturbance of the Earth magnetic field .

Plot

X-axis: data and time in UTC. Ranges from 7 days in the left (left) to now (right).
Y-axis: coloured bars of 1 hour wide, height= K Belgium.
Height of a bar represents a natural number: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9

Sensors

The calculation is based upon the measurements done by the Royal Meteorological Institute of Belgium based on the measurements done in Dourbes and Manhay, Belgium.

Provider

Royal Meteorological Institute of Belgium

Data location

http://ionosphere.meteo.be/geomagnetism/K_BEL/

Proton Flux Passport



Type

Graph, flux of solar protons versus time.

Use

To quantify proton events impacting Earth.

Units

Particles/(cm^2*s*sr) with s=second and sr=steradian

Data location

https://www.swpc.noaa.gov/products/goes-proton-flux

Plot

X-axis: time in UTC
Y-axis: Particles/(cm^2*s*sr), i.e. how many solar protons with a particular energy (eV) pass in 1 sec through 1 square centimeter from a cone with the Sun at the apex. The count is done at geostationary height.
The 4 colours represent 4 energies.

Sensor

GOES-18 s

Sensor

Geostation

Caden

Integrated

SWEC
Space Weather Education Center