

# SPACE WEATHER INTRODUCTORY COURSE



Collaboration of




Solar-Terrestrial Centre of Excellence



Koninklijke luchtmacht



Koninklijk Nederlands  
Meteorologisch Instituut  
*Ministerie van Infrastructuur en Milieu*



## Sensors

Jan Janssens, Dr Christophe Marqué

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



H-alpha picture: [https://www.windows2universe.org/spaceweather/ESF\\_loop.html](https://www.windows2universe.org/spaceweather/ESF_loop.html) (NSO/Sacramento Peak)

# Contents


- **Groundbased sensors**

- Visible light
- Radio domain
  - Humain
- Magnetosphere-Ionosphere
- Geomagnetism
- Neutron monitors
  - Dourbes

- **Spacebased sensors**

- GOES
- SDO
- PROBA2
- SOHO
- ACE
- DSCOVR
- STEREO

- Tools
- Overviews



### **Groundbased sensors**

Jan Janssens, Dr Christophe Marqué


SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



H-alpha picture: [https://www.windows2universe.org/spaceweather/ESF\\_loop.html](https://www.windows2universe.org/spaceweather/ESF_loop.html) (NSO/Sacramento Peak)

# Visible light

- GONG Network
  - White Light
  - H-alpha
  - Magnetogram
- Sunspot number
  - SILSO
- USET
  - WL, Ha, CaIIK
  - 250 obs. days / yr
- Catania
- NOAA / SOON
- K-Cor



GONG: **Global Oscillation Network Group** (<http://gong.nso.edu/>)

Originally developed to study solar oscillations

6 observing stations worldwide observing the Sun 24/7

**The six sites comprising the GONG Network are:**

The Big Bear Solar Observatory in California, USA.

The High Altitude Observatory at Mauna Loa in Hawaii, USA.

The Learmonth Solar Observatory in Western Australia.

The Udaipur Solar Observatory in India.

The Observatorio del Teide in the Canary Islands.

The Cerro Tololo Interamerican Observatory in Chile.

## Links

H-alpha: <http://halph.nso.edu/index.html>


White Light: <https://gong2.nso.edu/products/mainView/table.php?configFile=configs/mainView.cfg>

Magnetogram: <https://gong2.nso.edu/products/mainView/table.php?configFile=configs/mainView.cfg>

# Visible light

- GONG Network
  - White Light
  - H-alpha
  - Magnetogram
- Sunspot number
  - SILSO
- USET
  - WL, H $\alpha$ , CaIIK
  - 250 obs. days / yr
- Catania
- NOAA / SOON
- K-Cor

SWx Collaboration between STCE, Koninklij



## Links

SILSO: <http://sidc.oma.be/silso/> (Sunspot Index and Long-term Solar Observations)

USET: <http://www.sidc.be/uset/> (Uccle Solar Equatorial Table)

Catania: <http://web.ct.astro.it/sun/draw.jpg>

Catania and NOAA data are used as input for SIDC SWx forecasting

From the solar event listing at <ftp://ftp.swpc.noaa.gov/pub/indices/events/README>

Obs - The reporting observatory.

CUL - Culgoora, Australia

HOL - Holloman AFB, NM, USA

LEA - Learmonth, Australia

PAL - Palahua, HI, USA

RAM - Ramey AFB, PR, USA

SAG - Sagamore Hill, MA, USA

SVI - San Vito, Italy

Events from GOES satellites data show the SWPC Primary or Secondary GOES spacecraft for the observatory, e.g. G12

SOON: [https://en.wikipedia.org/wiki/Solar\\_Observing\\_Optical\\_Network](https://en.wikipedia.org/wiki/Solar_Observing_Optical_Network)

The SOON observatories are operated by detachments of AFWA's 2nd Weather Group at the following sites: RAAF Learmonth, Western Australia, Australia

Holloman AFB, New Mexico, USA

San Vito dei Normanni Air Station, San Vito dei Normanni, Italy (contractor-run site)

**Telescopes at Palahua, Hawaii and Ramey Air Force Base, Puerto Rico have been shut down.**

ISOON: <http://nsosp.nso.edu/isoon>


The planned **Improved Solar Observing Optical Network (ISOON)** is intended to replace the current SOON network. As of 2012, ISOON only exists at a single pilot site on Kirtland Air Force Base. *Images will be available when the telescope is working again.*

K-cor: <http://download.hao.ucar.edu/d5/www/fullres/latest/latest.kcor.gif>

Provides coronagraphic imagery. Large data gaps

# Sunspot number & Solar cycle

- Sunspot number
  - Sn
    - Also called International Sunspot Number (ISN)
  - $Sn = 10.g + s$ ,
    - with g the number of groups, and s the number of spots
  - Determined by the SIDC/SILSO (Uccle)!
    - <http://www.sidc.be/silso>
    - Network of about 80 stations



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



## Links

SILSO: <http://sidc.oma.be/silso/> (Sunspot Index and Long-term Solar Observations)

USET: <http://www.sidc.be/uset/> (Uccle Solar Equatorial Table)

Catania: <http://web.ct.astro.it/sun/draw.jpg>

The International sunspot number is a quantity that measures the number of sunspots and groups of sunspots present on the surface of the sun.

It is computed from a number of international observers using the formula:

$$R = k ( 10 g + s )$$

where


s is the number of individual spots,

g is the number of sunspot groups, and


k is a factor that varies with location and instrumentation (also known as the observatory factor or the personal reduction coefficient). It is not to be computed or applied by the observer.

# Sunspot number & Solar cycle

- Solar cycle
  - Avg. duration: +/- 11 years
    - Rise/Fall time: +/- 4 & 7 years
  - Avg.  $S_{n_{max}}$ : 184 (+/- 59)
    - SC24
      - Minimum: December 2008
      - Maximum: April 2014
      - $S_{n_{max}} = 116.4$
  - Butterfly diagram
    - Spots first appear at moderate latitudes (+/- 30°), then gradually move to equator
    - During SC minimum, groups of old and new SC exist



SILSO graphics (<http://sidc.be/silso>) Royal Observatory of Belgium 2019 March 1



SWIC – Collaboration between STCE, Koninklijke Luchtm

<http://www.stce.be/news/414/welcome.html>

## Links

SILSO: <http://sidc.oma.be/silso/> (Sunspot Index and Long-term Solar Observations)

USET: <http://www.sidc.be/uset/> (Uccle Solar Equatorial Table)

Catania: <http://web.ct.astro.it/sun/draw.jpg>

The International sunspot number is a quantity that measures the number of sunspots and groups of sunspots present on the surface of the sun.

It is computed from a number of international observers using the formula:

$$R = k ( 10 g + s )$$

where


s is the number of individual spots,

g is the number of sunspot groups, and

k is a factor that varies with location and instrumentation (also known as the observatory factor or the personal reduction coefficient). It is not to be computed or applied by the observer.



# Ground views from the Sun



SWIC = Collaboration between STCE, Koninklijke Luchtmacht, KNMI



## Links

USET: <http://www.sidc.be/uset/> (Uccle Solar Equatorial Table)

NSO GONG Magnetograms:

<https://gong2.nso.edu/products/tableView/table.php?configFile=configs/averageMagnetogram10min.cfg>

NSO GONG H-alpha: <http://halpha.nso.edu/>

Top left: USET: white light (rotated)

Top right: GONG: magnetogram

Bottom right: GONG: H-alpha (656.3 nm)

Bottom left: USET: Ca K (393.4 nm) – (rotated)

White Light: sunspots

Magnetogram: magnetic properties of sunspot groups and whole disk

H-alpha: filaments/prominences, flares

Ca K: plages (solar radiation)

NOAA/SWPC glossary at <https://www.swpc.noaa.gov/content/space-weather-glossary>

**Sunspot:** An area seen as a dark spot, in contrast with its surroundings, on the photosphere of the Sun. Sunspots are concentrations of magnetic flux, typically occurring in bipolar clusters or groups. They appear dark because they are cooler than the surrounding photosphere. Larger and darker sunspots sometimes are surrounded (completely or partially) by penumbrae. The dark centers are umbrae. The smallest, immature spots are sometimes called pores.

**Filaments/Prominences:** A mass of gas suspended over the chromosphere by magnetic fields and seen as dark ribbons threaded over the solar disk. A filament on the limb of the Sun seen in emission against the dark sky is called a prominence.

**Plages:** A brighter, hotter patch in the Sun's chromosphere, visible in H-alpha light and the calcium K line. Plages are the chromospheric equivalent of faculae on the photosphere, as can be seen when an active region is near the limb. Faculae have a strong influence on the solar constant, and the more readily detectable (because chromospheric) plage areas traditionally are used to monitor this influence.

Also good explanations at NASA/MSFC: <https://solarscience.msfc.nasa.gov/feature2.shtml> and at <https://astronomyconnect.com/forums/articles/7-observing-the-sun-in-ca-k-ca-h-and-other-narrow-bandwidths.38/>

- SWH Collaboration between STCE, Koninklijke Luchtmacht, KNMI

[illegible]

Catania info (Last update: 2017-Jan-24)						NOAA info (Last update: 2017-Jan-24)					Probabilities for			
Number	area	isots	Zurich	Longitude	Latitude	Number	MacIntosh	Mag. type	Longitude	Latitude	C flare	M flare	X flare	Proton
78	1	2	A	66.0	7.0	2626	H9x	Alpha	63.0	8.0	--	--	--	--
80	3	7	C	16.0	6.0	2627	D4L	Beta	12.0	6.0	--	--	--	--
B1	19	14	D	-2.0	12.0	2628	D5o	Beta	-7.0	12.0	--	--	--	--

10



Catania: <http://web.ct.astro.it/sun/draw.jpg>

Catania and NOAA data are used as input for SIDC SWx forecasting

:Issued: 2014 Apr 17 1325 UTC  
:Product: documentation at <http://www.sidc.be/products/tot>  
#-----#  
# DAILY BULLETIN ON SOLAR AND GEOMAGNETIC ACTIVITY from the SIDC #  
#-----#  
SIDC URSIGRAM 40417  
SIDC SOLAR BULLETIN 17 Apr 2014, 1304UT

SIDC FORECAST (valid from 1230UT, 17 Apr 2014 until 19 Apr 2014)  
SOLAR FLARES : Active (M-class flares expected, probability >=50%)  
GEOMAGNETISM : Quiet (A<20 and K<4)  
SOLAR PROTONS : Quiet

PREDICTIONS FOR 17 Apr 2014 10CM FLUX: 180 / AP: 013  
PREDICTIONS FOR 18 Apr 2014 10CM FLUX: 184 / AP: 007  
PREDICTIONS FOR 19 Apr 2014 10CM FLUX: 188 / AP: 005

COMMENT: Eleven sunspot groups were reported by NOAA today. NOAA ARs 2035, 2036, and 2037 (Catania numbers 24, 25, and 26 respectively) maintain the beta-gamma configuration of the photospheric magnetic field. The strongest flare of the past 24 hours was the M1.0 flare peaking at 19:59 UT yesterday in the NOAA AR 2035 (Catania number 24). The flare was associated with an EIT wave and a weak coronal dimming, but the associated CME was narrow and is not expected to arrive at the Earth.

We expect further flaring activity on the C-level, especially in the NOAA ARs 2035 and 2037 (Catania numbers 24 and 26 respectively) as well as in the NOAA AR 2042 (no Catania number yet) that yesterday appeared from behind the east solar limb, with a good chance for an M-class event.

Since yesterday evening the Earth is situated inside a solar wind structure with an elevated interplanetary magnetic field magnitude (occasionally up to 10 nT). It may be a weak ICME or the compression region on the flank of an ICME that missed the Earth. The solar origin of this structure is not clear. The north-south magnetic field component Bz was not strong, so no significant geomagnetic disturbance resulted (K index stayed below 4). Currently the solar wind speed is around 380 km/s and the IMF magnitude is around 8 nT.

We expect quiet to unsettled (K index up to 3) geomagnetic conditions, with active geomagnetic conditions (K = 4) possible, but unlikely.

TODAY'S ESTIMATED ISN : 145, BASED ON 17 STATIONS.  
99999

SOLAR INDICES FOR 16 Apr 2014  
WOLF NUMBER CATANIA : ///  
10CM SOLAR FLUX : 184  
AK CHAMBON LA FORET : 012  
AK WINGST : 004  
ESTIMATED AP : 004  
ESTIMATED ISN : 139, BASED ON 29 STATIONS.

NOTICEABLE EVENTS SUMMARY  
DAY BEGIN MAX END LOC XRAY OP 10CM Catania/ NOAA RADIO\_BURST\_TYPES  
16 1954 1959 2004 S14E09 M1.0 1N 24/2035 II/2  
END




*Finding your way  
in the  
URSIGram*

## Catania regions

## Sunspot numbers

# Visible light

- GONG Network
  - White Light
  - H-alpha
  - Magnetogram
- Sunspot number
  - SILSO
- USET
  - WL, Ha, CaIIK
  - 250 obs. days / yr
- Catania
- NOAA / SOON
- K-Cor



From the solar event listing at <ftp://ftp.swpc.noaa.gov/pub/indices/events/README>

Obs - The reporting observatory.

CUL - Culgoora, Australia

HOL - Holloman AFB, NM, USA

LEA - Learmonth, Australia

PAL - Palahua, HI, USA

RAM - Ramey AFB, PR, USA

SAG - Sagamore Hill, MA, USA

SVI - San Vito, Italy

Events from GOES satellites data show the SWPC Primary or Secondary GOES spacecraft for the observatory, e.g. G12

SOON: [https://en.wikipedia.org/wiki/Solar\\_Observing\\_Optical\\_Network](https://en.wikipedia.org/wiki/Solar_Observing_Optical_Network)

The SOON observatories are operated by detachments of AFWA's 2nd Weather Group at the following sites: RAAF Learmonth, Western Australia, Australia

Holloman AFB, New Mexico, USA

San Vito dei Normanni Air Station, San Vito dei Normanni, Italy (contractor-run site)

**Telescopes at Palahua, Hawaii and Ramey Air Force Base, Puerto Rico have been shut down.**

ISOON: <http://nsosp.nso.edu/isoon>

The planned **Improved Solar Observing Optical Network (ISOON)** was intended to replace the current SOON network. As of 2012, ISOON only exists at a single pilot site on Kirtland Air Force Base. *Images will be available when the telescope is working again.*

Giersch et al. (2018) - Reanalysis of Solar Observing Optical Network Sunspot Areas


<http://adsabs.harvard.edu/abs/2018SoPh..293..138G>

SOON began operations in the mid-1970s with four sites (Holloman, New Mexico, USA; Ramey, Puerto Rico; Learmonth, Western Australia; and Palahua, Hawaii, USA) operating by 1980. In 1987, San Vito, Italy, became operational. ... In 1996, the Palahua SOON equipment was returned to the National Solar Observatory at Sacramento Peak, New Mexico, to be used in the development of the Improved Solar Observing Optical Network (ISOON). However, **ISOON has never been deployed**. The Ramey site was closed in 2002. Thus, currently, there are only three SOON sites operating.

K-cor: <http://download.hao.ucar.edu/d5/www/fullres/latest/latest.kcor.gif>

Also at <https://www2.hao.ucar.edu/cosmo/k-cor>

Provides coronagraphic imagery. Large data gaps, limited time coverage during the day.



The flux at 2800 MHz (10.7 cm) is measured since 1946-1947 in Canada. First in Ottawa, then in 1962 in Algonquin Radio observatory, 250 km away from Ottawa, and since 1990 in Pentiction. This is the longest time series reflecting the solar activity besides the Sunspot Index (to which it highly correlates). More info in : K. F. Tapping, "The 10.7 cm solar radio flux (F10.7)", Space Weather, 11, 394, 2013

DRAO: Dominion Radio Astrophysical Observatory  
 (<https://www.nrc-cnrc.gc.ca/eng/solutions/facilities/drao.html> )

The Radio Solar Telescope Network (RSTN) started its operation during the 1970s and is operated by the US Air Force. It consists of an ensemble of flux monitoring instruments and a set of radio spectrometers. Data are not available in real time, but reports of events are provided through NOAA. Data become available for scientists several months (or years!) afterwards.

# NOAA radio event list

Event #	Time Start Max End	Station	Burst RBR, RSP	Frequency/band	Magnitude / type
2320 +	1155 1203	1206 G15 5	XRA 1-8A	M2.5	7.3E-03 2445
2320 +	1200 1200	1202 SAG G	RBR 1415	34	2445
2320 +	1201 1202	1203 SVI G	RBR 8800	94	2445
2320 +	1201 1202	1203 SAG G	RBR 610	750	2445
2320 +	1201 1202	1203 SVI G	RBR 4995	69	2445
2320 +	1202 1202	1203 SVI G	RBR 15400	66	2445
2320 +	1202 1202	1203 SAG G	RBR 2695	28	2445
2320 +	1202 1204	1205 SVI G	RBR 245	3600	2445
2320 +	1202 1204	1205 SVI G	RBR 410	047-171	2445
2320 +	1202 1204	1205 SVI G	RBR N12W73	2445	2445
2340 +	13127 U1339 A1348	SVI 2	FLA N09W04	2B	ERU 2443
2340 +	1331 1352	G15 5	XRA 1-8A	M3.7	5.9E-02 2443
2340 +	1336 1341	1413 SVI G	RBR 4995	740	2443
2340 +	1337 1341	1442 SVI G	RBR 2695	340	2443
2340 +	1337 1341	1429 SVI G	RBR 8800	560	2443
2340 +	1338 1341	1414 SVI G	RBR 15400	210	2443
2340 +	1343 1358	SAG C	RSP 048-180	11/2	955 2443
2340 +	1351 1531	SVI C	RSP 025-171	1V/1	2443
2340 +	1404 1426	1502 SAG G	RBR 410	1400	2443
2340 +	1405 1433	1507 SAG G	RBR 245	1400	2443
2340 +	1406 1427	1456 SAG G	RBR 1415	5800	2443
2340 +	1406 1427	1458 SAG G	RBR 610	1000	2443
2390	1421 1425	1433 SAG G	RBR 2695	180	

SWIC – Collaboration between STC, Koninklijke Luchtmacht, KNMI

14



NOAA radio event list available at <https://www.swpc.noaa.gov/products/solar-and-geophysical-event-reports>

Station Abbreviations:

SVI: San Vito

SAG: Sagamore Hill

PAL: Palehua

LEA: Learmonth

PEN: Penticton

Type of Emission:

RBR: Radio Burst at fixed frequency

RSP: Radio Burst identified by its type in spectral data (radio sweep)

Frequency:

Frequency of the burst (in MHz) or frequency range in which it is observed

Magnitude/type:

For bursts at fixed frequency: magnitude above quiet Sun in Solar Flux Unit

For bursts reported by type: type/magnitude (1-3: weak to strong). If type II a speed is given in km/s (here 955 km/s)

# Radio burst magnitudes

## Typ. Quiet Sun values [SFU]

Frequency	Solar min.	Solar max. (Z=200)
245 MHz	10	15
410 MHz	25	35
610 MHz	30	45
1415 MHz	50	100
2695 MHz	70	200
2800 MHz	70	200
4995 MHz	100	200
8800 MHz	220	290
15400 MHz	580	650

How frequently bursts of a certain magnitude occur? 1 event every X days

f < 2000 MHz

Magnitude	Solar min.	Solar max.
1000	5 days	0.7 day
10000	34 days	4 days
100000	212 days	17 days

f > 2000 MHz

Magnitude	Solar min.	Solar max.
1000	38 days	6 days
10000	247 days	39 days
100000	1594 days	255 days

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



The left table reports typical values observed for the quiet Sun during minimum and maximum conditions. There can be lower and higher values but this is just to give an idea.

Values from the tables to the Right are derived from the paper by Nita et al. 2002. This gives in how many days on average a burst of a certain magnitude happens in each band. If we take a finer frequency band definition, the numbers can be different (see next slide).

Nita et al., 2002

The Peak Flux Distribution of Solar Radio Bursts

<http://adsabs.harvard.edu/abs/2002ApJ...570..423N>

# Radio burst magnitude

## 100 MHz < f < 900 MHz

Magnitude	Solar Min.	Solar Max.
1000	6 days	0.75 days
10000	36 days	4 days
100000	223 days	17 days

## 1000 MHz < f < 1700 MHz

Magnitude	Solar Min.	Solar Max.
1000	65 days	12 days
10000	385 days	75 days
100000	2266 days	450 days

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



The right table shows more interesting statistics for a frequency band that covers GNSS services and air traffic radar and surveillance bands



# Penticton

- Flux measurement at 2800 MHz (10.7 cm), 100 MHz bandwidth
- 3 times per day
- “official” value for the day is the one of 20:00 UT (local noon)
- Accuracy:
  - < 100 sfu: 1 sfu
  - > 100 sfu: 1% of flux
- Uncorrected for solar flares
- R-, S-, Q-component



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



Information from: K. F. Tapping, “The 10.7 cm solar radio flux (F10.7)”, Space Weather, 11, 394, 2013  
<http://adsabs.harvard.edu/abs/2013SpWea..11..394T>

Daily 10.7cm solar radio fluxes from Penticton at <http://www.spaceweather.ca/solarflux/sx-4a-en.php>

Daily radio fluxes (other wavelengths) at <http://legacy-www.swpc.noaa.gov/ftplib/lists/radio/rad.txt>

Solar flux unit:  
 1 sfu =  $10^{-22}$  W m<sup>-2</sup> Hz<sup>-1</sup>

The Humain solar radio observatory efforts to develop 10.7cm radio flux measurements were halted after it became clear there was too much interference from Military instruments. <http://www.sidc.be/humain/index.php>

From Tapping (2013):

[7] The early measurements of solar centimetric emissions were made using relatively small antennas, having beams subtending solid angles larger than that subtended by the solar disk, so no determinations of the distribution of emission could be made on any routine basis. These spatially integrated emissions were categorized on the basis of their characteristic timescale of variation into three identifiable components: a rapidly varying or R component, comprising emissions varying over timescales in the second-minute range, perhaps as long as an hour. Slower variations were lumped into a slowly varying or S component. Extrapolation to zero activity suggested an underlying constant, base level, which became called the quiet sun, or Q component. The terms R and Q have fallen out of use, and these components are now known, respectively, as bursts and the quiet sun background emission. The slowly varying component originates primarily in active regions; its intensity is a measure of the overall level of solar magnetic activity and has a broad spectral peak at about 10 cm wavelength. The F10.7 values comprise contributions from the S component and the quiet sun background, and sometimes from radio bursts.

:Issued: 2014 Apr 17 1325 UTC  
:Product: documentation at <http://www.sidc.be/products/tot>  
#-----#  
# DAILY BULLETIN ON SOLAR AND GEOMAGNETIC ACTIVITY from the SIDC #  
#-----#  
SIDC URSIGRAM 40417  
SIDC SOLAR BULLETIN 17 Apr 2014, 1304UT

SIDC FORECAST (valid from 1230UT, 17 Apr 2014 until 19 Apr 2014)  
SOLAR FLARES : Active (M-class flares expected, probability >=50%)  
GEOMAGNETISM : Quiet (A<20 and K<4)  
SOLAR PROTONS : Quiet

PREDICTIONS FOR 17 Apr 2014 10CM FLUX: 180 / AP: 013  
PREDICTIONS FOR 18 Apr 2014 10CM FLUX: 184 / AP: 007  
PREDICTIONS FOR 19 Apr 2014 10CM FLUX: 188 / AP: 005

COMMENT: Eleven sunspot groups were reported by NOAA today. NOAA ARs 2035, 2036, and 2037 (Catania numbers 24, 25, and 26 respectively) maintain the beta-gamma configuration of the photospheric magnetic field. The strongest flare of the past 24 hours was the M1.0 flare peaking at 19:59 UT yesterday in the NOAA AR 2035 (Catania number 24). The flare was associated with an EIT wave and a weak coronal dimming, but the associated CME was narrow and is not expected to arrive at the Earth.

We expect further flaring activity on the C-level, especially in the NOAA ARs 2035 and 2037 (Catania numbers 24 and 26 respectively) as well as in the NOAA AR 2042 (no Catania number yet) that yesterday appeared from behind the east solar limb, with a good chance for an M-class event.

Since yesterday evening the Earth is situated inside a solar wind structure with an elevated interplanetary magnetic field magnitude (occasionally up to 10 nT). It may be a weak ICME or the compression region on the flank of an ICME that missed the Earth. The solar origin of this structure is not clear. The north-south magnetic field component Bz was not strong, so no significant geomagnetic disturbance resulted (K index stayed below 4). Currently the solar wind speed is around 380 km/s and the IMF magnitude is around 8 nT.

We expect quiet to unsettled (K index up to 3) geomagnetic conditions, with active geomagnetic conditions (K = 4) possible, but unlikely.

TODAY'S ESTIMATED ISN : 145, BASED ON 17 STATIONS.  
99999

SOLAR INDICES FOR 16 Apr 2014

WOLF NUMBER CATANIA : -///

10CM SOLAR FLUX : 184

AK CHAMBON LA FORET : 012

AK WINGST : 004

ESTIMATED AP : 004

ESTIMATED ISN : 139, BASED ON 29 STATIONS.

NOTICEABLE EVENTS SUMMARY

DAY BEGIN MAX END LOC XRAY OP 10CM Catania/NOAA RADIO\_BURST\_TYPES

16 1954 1959 2004 S14E09 M1.0 1N 24/2035 II/2

END



*Finding your way  
in the  
URSIGram*


**10.7cm Radio flux**

# Nançay radio observations


- Nançay Radioheliograph
  - Imaging at several frequencies 150 – 450 MHz
- ORFEES spectrograph (130 – 1000 MHz)

<https://realtime.obs-nancay.fr/orfees/>

Archives: <http://secchirh.obspm.fr>



Luchtmacht, KN



The Nançay Radioheliograph (NRH) provides 2d imaging of the solar corona between 150 and 450 MHz. It reveals radio emission associated with active regions (with sunspots) called noise storms. When an eruptive event occurs, imaging of different types of bursts (type II, type III, type IV etc...) is made. When solar activity is low, the quiescent corona becomes visible, showing the extent of large scale coronal structures (streamers, coronal holes etc...). The instrument is currently off for hardware upgrade.

The ORFEES spectrograph covers the band 130 – 1000 MHz and complements the imaging observations by providing the spectral type of the bursts. It's an instrument specifically built for space weather operations by a joint effort between the Paris Observatory and the French Air Force.

ORFEES (Observation Radio Fréquences pour l'Etude des Eruptions Solaires)

<https://realtime.obs-nancay.fr/orfees/>


<https://www.obs-nancay.fr/-ORFEES-38-.html?lang=en>

The 17 July 2012 event:

1260 +	1203	1715	1904	G15	5	XRA	1-8A	M1.7	2.1E-01	1520
1330	1307	////	1952	SAG	C	RSP	039-180	CTM/1		
1260 +	1328	U1643	A1802	COM	3	FLA	S28W65	1F		1520
1280 +	1328	1348	1356	SVI	G	RNS	245	260		
1240	B1348	////	1730	SOH	4	CME	XUV,EUV,UV153-304/FS814			1520

# Callisto Network


- Network of identical low cost analog receivers (~130 stations over the world)
- <http://www.e-callisto.org/>
- **Real time in Europe**
  - Birr Castle (Ireland): BIR
  - Humain (Belgium): HUMAIN
  - Bleien (Switzerland): BLEN5M, BLEN5W
  - Trieste (Italy): TRIEST
  - Metsähovi (Finland): MRO
  - Kellyville (Greenland): GREENLAND



<http://soleil.i4ds.ch/solarradio/callistoQuicklooks/>

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

20

- # Callisto Network
- Network of identical low cost analog receivers (~130 stations over the world)
  - <http://www.e-callisto.org/>
  - **Real time in Europe**
    - Birr Castle (Ireland): BIR
    - Humain (Belgium): HUMAIN
    - Bleien (Switzerland): BLEN5M, BLEN5W
    - Trieste (Italy): TRIEST
    - Metsähovi (Finland): MRO
    - Kellyville (Greenland): GREENLAND
- 
- <http://soleil.i4ds.ch/solarradio/callistoQuicklooks/>
- SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI
- 20

Callisto stands for: Compound Astronomical Low cost Low frequency Instrument for Spectroscopy and Transportable Observatory  
e-callisto stands for the network of callisto observatories.

## European solar radio observatories *with real-time access*



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI


21



This map shows the locations of solar radio instruments operated by professional bodies (institutes, observatories etc...). Their data (essentially dynamic spectra) are available in real time or near real time

Available at e-Callisto at <http://soleil.i4ds.ch/solarradio/callistoQuicklooks/>

# The Humain station



About 120 km south-east of Brussels

# Humain: Solar instruments

- 6-m dish
  - Automated operations, Sun tracking ~7h30 – 16h00 UT
  - VHF antenna (piggy back)
  - UHF antenna at focus
- 
- VHF antenna (45 – 450 MHz)
    - Callisto receiver
    - ARCAS receiver
  - UHF antenna (275 – 1495 MHz)
    - HSRS receiver



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



The ARCAS and HSRS receivers are based on commercial Software Defined Radio receivers. The RF signal is digitized and all operations needed to create the dynamic spectrum is made by programming on a control PC. The development of ARCAS and HSRS was made at ROB.

Callisto stands for: Compound Astronomical Low cost Low frequency Instrument for Spectroscopy and Transportable Observatory

ARCAS stands for Augmented Resolution Callisto Spectrometer -

<http://www.stce.be/news/369/welcome.html>

HSRS stands for HUMAIN Solar Radio Spectrograph - <http://www.stce.be/news/326/welcome.html>

# Humain: Solar instruments

- 6-m dish
- Automated operations, Sun tracking ~7h30 – 16h00 UT
- VHF antenna (piggy back)
- UHF antenna at focus

- VHF antenna (45 – 450 MHz)
  - Callisto receiver
  - ARCAS receiver
- UHF antenna (275 – 1495 MHz)
  - HSRS receiver



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



The ARCAS and HSRS receivers are based on commercial Software Defined Radio receivers. The RF signal is digitized and all operations needed to create the dynamic spectrum is made by programming on a control PC. The development of ARCAS and HSRS was made at ROB.

Callisto stands for: Compound Astronomical Low cost Low frequency Instrument for Spectroscopy and Transportable Observatory

ARCAS stands for Augmented Resolution Callisto Spectrometer -

<http://www.stce.be/news/369/welcome.html>


HSRS stands for HUMAIN Solar Radio Spectrograph - <http://www.stce.be/news/326/welcome.html>



# Humain: Solar instruments

	Callisto	ARCAS	HSRS
Type	Analog receiver	Digital	Digital
Frequency band	45 – 447 MHz	45 – 450 MHz	275 – 1495 MHz
Frequency resolution	63 kHz	98 kHz	98 kHz
Time resolution	250 ms	~ 84 ms	~ 250 ms
# of frequencies	200	~ 4.2 k	~ 12.5 k

Data available in near realtime  
<http://sidc.be/humain>



See also the news item at <http://www.stce.be/news/384/welcome.html> for this event.

<https://www.cv.nrao.edu/course/astr534/Introradastro.html>

The Earth's ionosphere prevents ground-based observations at frequencies below 10 MHz (wavelengths higher than 30 m).

:Issued: 2014 Apr 17 1325 UTC  
:Product: documentation at <http://www.sidc.be/products/tot>  
#-----#  
# DAILY BULLETIN ON SOLAR AND GEOMAGNETIC ACTIVITY from the SIDC #  
#-----#  
SIDC URSIGRAM 40417  
SIDC SOLAR BULLETIN 17 Apr 2014, 1304UT

SIDC FORECAST (valid from 1230UT, 17 Apr 2014 until 19 Apr 2014)  
SOLAR FLARES : Active (M-class flares expected, probability >=50%)  
GEOMAGNETISM : Quiet (A<20 and K<4)  
SOLAR PROTONS : Quiet

PREDICTIONS FOR 17 Apr 2014 10CM FLUX: 180 / AP: 013  
PREDICTIONS FOR 18 Apr 2014 10CM FLUX: 184 / AP: 007  
PREDICTIONS FOR 19 Apr 2014 10CM FLUX: 188 / AP: 005

COMMENT: Eleven sunspot groups were reported by NOAA today. NOAA ARs 2035, 2036, and 2037 (Catania numbers 24, 25, and 26 respectively) maintain the beta-gamma configuration of the photospheric magnetic field. The strongest flare of the past 24 hours was the M1.0 flare peaking at 19:59 UT yesterday in the NOAA AR 2035 (Catania number 24). The flare was associated with an EIT wave and a weak coronal dimming, but the associated CME was narrow and is not expected to arrive at the Earth.

We expect further flaring activity on the C-level, especially in the NOAA ARs 2035 and 2037 (Catania numbers 24 and 26 respectively) as well as in the NOAA AR 2042 (no Catania number yet) that yesterday appeared from behind the east solar limb, with a good chance for an M-class event.

Since yesterday evening the Earth is situated inside a solar wind structure with an elevated interplanetary magnetic field magnitude (occasionally up to 10 nT). It may be a weak ICME or the compression region on the flank of an ICME that missed the Earth. The solar origin of this structure is not clear. The north-south magnetic field component Bz was not strong, so no significant geomagnetic disturbance resulted (K index stayed below 4). Currently the solar wind speed is around 380 km/s and the IMF magnitude is around 8 nT.

We expect quiet to unsettled (K index up to 3) geomagnetic conditions, with active geomagnetic conditions (K = 4) possible, but unlikely.

TODAY'S ESTIMATED ISN : 145, BASED ON 17 STATIONS.  
99999

SOLAR INDICES FOR 16 Apr 2014  
WOLF NUMBER CATANIA : ///  
10CM SOLAR FLUX : 184  
AK CHAMBON LA FORET : 012  
AK WINGST : 004  
ESTIMATED AP : 004  
ESTIMATED ISN : 139, BASED ON 29 STATIONS.

#### NOTICEABLE EVENTS SUMMARY

DAY	BEGIN	MAX	END	LOC	XRAY	OP	10CM	Catania/NOAA	RADIO_BURST_TYPES
16	1954	1959	2004	S14E09	M1.0	IN		24/2035	II/2
END									



*Finding your way  
in the  
URSIGram*

*Radio bursts*

# Magnetosphere - Ionosphere

## Magnetosphere

- Magnetometers
- Neutron monitors
- ...

– See [Earth Environment - Magnetosphere](#)




## Ionosphere

- Ionospheric sounders
- GNSS
- ...

– See [Earth Environment – Ionosphere](#)

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

# Geomagnetic indices



SWPC Kp index: <https://www.swpc.noaa.gov/products/planetary-k-index>

Dst index (Kyoto WDC): [http://wdc.kugi.kyoto-u.ac.jp/dst\\_realtime/presentmonth/index.html](http://wdc.kugi.kyoto-u.ac.jp/dst_realtime/presentmonth/index.html)

GOES Hp: <https://www.swpc.noaa.gov/products/goes-magnetometer>

<https://www.swpc.noaa.gov/sites/default/files/images/u2/TheK-index.pdf>

The A-index was invented because there was a need to derive some kind of daily average level for geomagnetic activity. Because of the non-linear relationship of the K-scale to magnetometer fluctuations, it is not meaningful to take averages of a set of K indices.

<http://www.stce.be/news/243/welcome.html>

<http://www.stce.be/news/301/welcome.html>


Cander et al. (1998): Forecasting ionospheric structure during the great geomagnetic storms

<http://adsabs.harvard.edu/abs/1998JGR...103..391C>

The size of a geomagnetic storm is classified as moderate ( $-50 \text{ nT} > \text{minimum of Dst} > -100 \text{ nT}$ ), intense ( $-100 \text{ nT} > \text{minimum Dst} > -250 \text{ nT}$ ) or super-storm (minimum of Dst  $< -250 \text{ nT}$ ).

# Dourbes

K-type index of local magnetic activity, Dourbes (50.1°N, 4.6°E)  
(copyright RMI)



Dourbes:

Geomagnetism: [http://ionosphere.meteo.be/geomagnetism/ground\\_K\\_dourbes](http://ionosphere.meteo.be/geomagnetism/ground_K_dourbes)

Neutron monitor: <http://www.nmdb.eu/>

Ionosphere: <http://digisonde.oma.be/>

## Exercise: 10.7cm Radio Flux

- It is 17 April 2014, and the solar cycle (SC24) is in its period of maximum solar activity. The 10.7cm radio flux for that day (at 20UT) is reported to be 600 sfu. This is:
  - A typical value for the radio flux during SC max
  - A radio flux value affected by a strong solar flare

# Contents

- **Groundbased sensors**


- Visible light
- Radio domain
  - Humain
- Magnetosphere-Ionosphere
- Geomagnetism
- Neutron monitors
  - Dourbes

- **Spacebased sensors**

- GOES
- SDO
- PROBA2
- SOHO
- ACE
- DSCOVR
- STEREO

- Tools
- Overviews





## Spacebased sensors

Jan Janssens


SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

33 


PROBA2 and picture of the Sun

# Why do we need SWx satellites?

- EUV and X-ray (solar atmosphere)
  - Flares & Coronal holes
- Coronagraphs
- Solar wind (in-situ)
- Solar farside
  - 20 September 2012
  - 23 July 2012
  - ...
- Radio
  - Triangulation
  - Low frequencies
- Science
- White light (24hrs)
- ... SWIC – Collaboration between STCE, Konink



More on the 20 September 2012 flare at <http://www.stce.be/news/263/welcome.html>



Lagrangian points: [https://en.wikipedia.org/wiki/Lagrangian\\_point](https://en.wikipedia.org/wiki/Lagrangian_point)

Earth orbits: [https://en.wikipedia.org/wiki/List\\_of\\_orbits#Altitude\\_classifications\\_for\\_geocentric\\_orbits](https://en.wikipedia.org/wiki/List_of_orbits#Altitude_classifications_for_geocentric_orbits)

\* GEO: GOES, SDO (inclined)

Advantages and disadvantages of SDO in GEO at <https://sdo.gsfc.nasa.gov/mission/project.php>


### Orbit

The rapid cadence and continuous coverage required for SDO observations led to placing the satellite into an inclined geosynchronous orbit. This allows for a nearly-continuous, high-data-rate, contact with a single, dedicated, ground station.

Nearly continuous observations of the Sun can be obtained from other orbits, such as low Earth orbit (LEO). If SDO were placed into an LEO it would be necessary to store large volumes of scientific data onboard until a downlink opportunity. The large data rate of SDO, along with the difficulties in managing a large on-board storage system, resulted in a requirement of continuous contact.

The disadvantages of this orbit include higher launch and orbit acquisition costs (relative to LEO) and eclipse (Earth shadow) seasons twice annually. During these 2-3 week eclipse periods, SDO will experience a daily interruption of solar observations. There will also be three lunar shadow events each year from this orbit.

This orbit is located on the outer reaches of the Earth's radiation belt where the radiation dose can be quite high. Additional shielding was added to the instruments and electronics to reduce the problems caused by exposure to radiation. Because this is a Space Weather effect, SDO is affected by the very processes it is designed to study!



Lagrangian points: [https://en.wikipedia.org/wiki/Lagrangian\\_point](https://en.wikipedia.org/wiki/Lagrangian_point)

Earth orbits: [https://en.wikipedia.org/wiki/List\\_of\\_orbits#Altitude\\_classifications\\_for\\_geocentric\\_orbits](https://en.wikipedia.org/wiki/List_of_orbits#Altitude_classifications_for_geocentric_orbits)

\* M/HEO: **INTEGRAL** (INTERNational Gamma-Ray Astrophysics Laboratory)

[http://space-env.esa.int/index.php/SREM\\_Plots.html](http://space-env.esa.int/index.php/SREM_Plots.html)

SREM: Standard Radiation Environment Monitor

[http://srem.web.psi.ch/html/srem\\_home.shtml](http://srem.web.psi.ch/html/srem_home.shtml)

Integral is the last remaining operational radiation monitor.

\* LEO: PROBA2, HINODE, RHESSI, FERMI

**Hinode**: <http://hinode.nao.ac.jp/gallery/latest/>

XRT: X-Ray Telescope ; Also at <https://www.solarmonitor.org/>

**RHESSI**: Reuven Ramaty High Energy Solar Spectroscopic Imager

<https://hesperia.gsfc.nasa.gov/rhessi3/>

**PROBA2**: PProject for Onboard Autonomy

<http://proba2.oma.be/ssa>

**FERMI**: Fermi Gamma-ray Space Telescope

<http://www.astronomerstelegram.org/?read=10720> (detections of solar gamma ray bursts)

\* L1: First Lagrangian point

DISCOVER, ACE, SOHO


**Wind**: <https://pwg.gsfc.nasa.gov/windnrt/>

\* Solar orbit

STEREO

# GOES

- X-ray flux
- Proton flux
- Magnetic field
- Electron flux
- Imagery
  - GOES-12-15
    - X-ray: SXI
  - GOES-16
    - EUV: SUVI



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

37



**X-ray flux:** <https://www.swpc.noaa.gov/products/goes-x-ray-flux>

More info at [https://www.ngdc.noaa.gov/stp/satellite/goes/doc/GOES\\_XRS\\_readme.pdf](https://www.ngdc.noaa.gov/stp/satellite/goes/doc/GOES_XRS_readme.pdf)

**Proton flux:** <https://www.swpc.noaa.gov/products/goes-proton-flux>



**Magnetic field:** <https://www.swpc.noaa.gov/products/goes-magnetometer>

**Electron flux:** <https://www.swpc.noaa.gov/products/goes-electron-flux>

**Imagery (X-ray):** <https://www.swpc.noaa.gov/products/goes-solar-x-ray-imager-sxi>

# GOES

- X-ray flux
- Proton flux
- Magnetic field
- Electron flux
- Imagery
  - GOES-12-15
    - X-ray: SXI
      - Discontinued
  - GOES-16
    - EUV: SUVI
      - Operational



SWIC – Collaboration between STCE, Koninklijke

**Imagery (X-ray):** <https://www.swpc.noaa.gov/products/goes-solar-x-ray-imager-sxi>  
Solar X-ray Imager

GOES-16/SUVI has been operationally checked out.  
SUVI and 4 other instruments onboard GOES-16 have been put in safe mode as of 20 December 2017.  
Entering operation in December 2019, becoming primary solar imager on 9 December 2019.  
<https://www.swpc.noaa.gov/news/noaanesdis-has-extended-operational-period-both-goes-1415-02-march-2020>  
<https://www.swpc.noaa.gov/products/goes-solar-ultraviolet-imager-suvi>  
<https://www.nesdis.noaa.gov/GOES-R-Series-Satellites>

**Imagery (SUVI):** <https://www.goes-r.gov/spacesegment/suvi.html>  
Solar UltraViolet Imager

# SUVI


Wavelength Log (Te)	94 Å 6.8	131 Å 7.0,7.2	171 Å 5.8	195 Å 6.1,7.3	284 Å 6.3	304 Å 4.7
Filaments						
Coronal Holes						
Active Region Complexity						
CMEs (e.g. dimming)						
Flare Location and Morphology						
Quiet Regions						

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

Table from <https://www.goes-r.gov/spacesegment/suvi.html>

# SDO

- HMI
  - Helioseismic and Magnetic Imager
  - « White light » and Magnetograms
- AIA
  - Atmospheric Imaging Assembly
  - EUV imagery in 10 filters
- EVE
  - Extreme ultraviolet Variability Experiment
  - Scaled to GOES x-ray measurements



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

Imagery and data at <https://sdo.gsfc.nasa.gov/data/>


AIA: Instrument description and characteristics of filters:

Boerner et al., 2012: <http://adsabs.harvard.edu/abs/2012SoPh..275...41B>

[http://jsoc.stanford.edu/HMI/docs/AIA\\_calibration.pdf](http://jsoc.stanford.edu/HMI/docs/AIA_calibration.pdf)

[https://www.nasa.gov/pdf/417176main\\_SDO\\_Guide\\_CMR.pdf](https://www.nasa.gov/pdf/417176main_SDO_Guide_CMR.pdf)





ESP: EUV SpectroPhotometer

MEGS: multiple EUV grating spectrograph (-A no longer operational)

SAM: Solar Aspect Monitor (no longer operational)

The EVE proxy for x-ray flux is at [http://lasp.colorado.edu/eve/data\\_access/sdo-goes-eve-flare-watch/index.html](http://lasp.colorado.edu/eve/data_access/sdo-goes-eve-flare-watch/index.html)

B9.5 flare on 22 January 2018

[http://lasp.colorado.edu/eve/data\\_access/](http://lasp.colorado.edu/eve/data_access/)




Crotser et al. (2004): SDO-EVE multiple EUV grating spectrograph (MEGS) optical design

<http://adsabs.harvard.edu/abs/2004SPIE.5563..182C>

The NASA Solar Dynamics Observatory (SDO), scheduled for launch in 2008, incorporates a suite of instruments including the EUV Variability Experiment (EVE). The EVE instrument package contains grating spectrographs used to measure the solar extreme ultraviolet (EUV) irradiance from 0.1 to 105 nm. The Multiple EUV Grating Spectrograph (MEGS) channels use concave reflection gratings to image solar spectra onto CCDs that are operated at -100°C. MEGS provides 0.1nm spectral resolution between 5-105nm every 10 seconds with an absolute accuracy of better than 25% over the SDO 5-year mission. MEGS-A utilizes a unique grazing-incidence, off-Rowland circle (RC) design to minimize angle of incidence at the detector while meeting high resolution requirements. MEGS-B utilizes a double-pass, cross-dispersed double-Rowland circle design. MEGS-P, a Ly- $\alpha$  monitor, will provide a proxy model calibration in the 60-105 nm range. Finally, the Solar Aspect Monitor (SAM) channel will provide continual pointing information for EVE as well as low-resolution X-ray images of the sun. In-flight calibrations for MEGS will be provided by the on-board EUV Spectrophotometer (ESP) in the 0.1-7nm and 17-37nm ranges, as well as from annual under-flight rocket experiments. We present the methodology used to develop the MEGS optical design.

# SDO

- AIA
  - Some filters peak at multiple temperatures
  - AIA 4500 no longer in use



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

More info at <http://aia.lmsal.com/public/instrument.htm>  
 And at <https://www.nasa.gov/content/goddard/how-sdo-sees-the-sun>

# SDO/AIA: Filters & Features

Filter/Wavelength (Å)	1700 6K	1600 10K	304 80K	171 700K	193 1.25M	211 2M	335 2.8M	94 6M	131 10M
Coronal holes				\	X	X	\		
CMEs: Coronal wave/dimming				\	X	X			
CMEs: ejections			X	X	X	\			\
Coronal loops			\	X	\				\
Flare ribbons		X	\	\	\	\			
Flare locations		\					\	X	X
Flares				\	\	\	\	X	X
Jets			X	\			\	\	X
Filament channels				\	X	X			
Filaments/Prominences			X	\	X	X			\
Sunspots		X							

X : Optimal line to see feature; \ : Reasonably visible ; "blanc" denotes barely or not visible

\* : Most filters peak at more than 1 temperature, have a broad bandpass, and see many of the features mentioned.  
The 1700Å filter is contaminated by continuum and C IV.

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI




More info at <http://aia.lmsal.com/public/instrument.htm>

And at <https://www.nasa.gov/content/goddard/how-sdo-sees-the-sun>

Table in collaboration with Dr Matt West (ROB/SIDC).

# SDO



*Credits: NASA / Goddard Space Flight Centre*

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI




Credits: NASA/GSFC


<https://svs.gsfc.nasa.gov/cgi-bin/details.cgi?aid=11385>

# PROBA2

**SWAP**



**LYRA**



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

45




Data and imagery at <http://proba2.oma.be/ssa>

From P2SC: <http://p2web.oma.be/about/sciencePayload>

LYRA (**L**arge **Y**ield **R**adiometer, formerly **LY**man alpha **R**adiometer) is an ultraviolet irradiance radiometer that observes the Sun in four passbands, chosen for their relevance to solar physics, aeronomy and space weather. This instrument can also detect flares and analyze the atmospheric composition of the Earth.

SWAP (**S**un **W**atcher using **A**ctive Pixel System detector and Image **P**rocessing) is a small EUV telescope that images the solar corona with a bandpass around 17.4 nm, corresponding to a temperature of 1 million degrees. SWAP continues the systematic CME watch program of EIT at an improved cadence and monitors events in the lower solar corona that might be relevant for space weather.

# SOHO



SOHO: Solar and Heliospheric Observatory

Launched on 2 December 1995 and still observing from the L1 point!

EIT and coronagraphic imagery at <https://sohowww.nascom.nasa.gov/data/realtime-images.html>

The passbands and related temperatures are similar to SDO/AIA, with the 28.4 nm filter corresponding to 2 million degrees.

Solar wind data (and shocks) at <http://umtof.umd.edu/pm/>

EIT: Extreme ultraviolet Imaging Telescope

CELIAS: Charge, Element, and Isotope Analysis System

MTOF: Mass Tome-of-Flight sensor

LASCO: Large Angle and Spectrometric Coronagraph

<http://star.mpae.gwdg.de/>


- C2 (1.5-6 solar radii)
- C3 (outer: 3-32 solar radii)

The C1 (inner: 1.1-3 solar radii) is no longer operational following SOHO's summer holidays in 1998:

<https://lasco-www.nrl.navy.mil/index.php?p=content/intro>

LASCO comprises of three telescopes (C1, C2 and C3), each of which looks at an increasingly large area surrounding the Sun. For the first year-and-a-half of the SOHO mission, all three instruments worked perfectly. However, in 1998 SOHO was accidentally "lost" in space after it received a bad command. The entire spacecraft lost power and essentially froze solid for several weeks. Eventually -- miraculously! -- the SOHO team were able to relocate the spacecraft, regain control and slowly power-up and thaw out the instruments. Sadly, the LASCO C1 camera was lost as a result of this but the rest of spacecraft came through almost completely unscathed! Eighteen years later -- and over twenty years since launch -- LASCO C2 and C3 (and most of the rest of SOHO!) continue to work extremely well, sending back images and data on a daily basis.

# SOHO




The weak CMEs/structures become better visible in difference imagery (one image subtracted from the other). Movies for the last 4 days can be found at <https://sohowww.nascom.nasa.gov/data/realtime/mpeg/> (resp. LASCO C2 combo and LASCO C3 combo).

SOHO/LASCO is also a notorious comet chaser, discovering its 3000th comet on 14 September 2015.

SOHO has/had many other instruments, amongst which was the famous MDI (Michelson Doppler Imager) which made images of the Sun in “white light” as well as magnetograms. This program was terminated in 2011 as it was superseded by SDO/HMI. It was exceptionally switched on again for the Mercury transit 2016 (<https://soho.nascom.nasa.gov/pickoftheweek/>). The MDI archives are at <http://soi.stanford.edu/>

Descriptions of the other instruments can be found at <https://sohowww.nascom.nasa.gov/about/instruments.html>

# ACE



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

48

ACE: Advanced Composition Explorer  
<http://www.srl.caltech.edu/ACE/>

All data at NOAA/SWPC: <https://www.swpc.noaa.gov/products/ace-real-time-solar-wind>

EPAM: Electron Proton Alpha Monitor

- measurements of low-energy electrons and protons

SWEPAM: Solar Wind Electron Proton Alpha Monitor

- measurements of solar wind density, speed and temperature

MAG: Magnetometer instrument

- Measurements of solar wind magnetic field

SIS: Solar Isotope Spectrometer


- Measurements of high-energy protons (>10 MeV, > 30 MeV)

ACE measurements of e.g. solar wind speed are sometimes unreliable in case of strong proton events, which is annoying when a CME arrives.

As of 27 July 2016, ACE is superseded by the DSCOVR satellite, and as such there's no longer a 24hr data coverage anymore.



# DSCOVER



DSCOVER: Deep Space Climate Observatory

<https://www.nesdis.noaa.gov/content/dscovr-deep-space-climate-observatory>

Solar wind data are available at <https://www.swpc.noaa.gov/products/real-time-solar-wind>

From NOAA/SWPC: Real-Time Solar Wind (RTSW) data refers to data from any spacecraft located upwind of Earth, typically orbiting the L1 Lagrange point, that is being tracked by the Real-Time Solar Wind Network of tracking stations. The NOAA DSCOVER satellite became the operational RTSW spacecraft on July 27, 2016 at 1600UT (noon EDT, 10am MDT).

SWPC maintains the ability to instantaneously switch the spacecraft that provides the RTSW data. During times of outages in DSCOVER data or problems with the data, this page may instead display the data from the NASA/ACE spacecraft.



The two DSCOVER instruments for which data are available:

Faraday Cup (FC) of the Harvard Smithsonian Astrophysical Observatory(link is external)

Magnetometer (MAG) of the University of NASA Goddard Space Flight Center (link is external)

Note that DSCOVER is vulnerable to GCR, which introduces spurious measurement readings from time to time. The correctness of the values can be checked against the ACE data. More on this issue at <http://www.nature.com/news/cosmic-rays-may-threaten-space-weather-satellite-1.20880>  
DSCOVER was in safe mode from 27 June 2019 till 02 March 2020 due to a technical glitch.  
<https://www.nesdis.noaa.gov/content/noaas-dscovr-satellite-operating-again>

# STEREO



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



STEREO: Solar-Terrestrial Relations Observatory - SSC: STEREO Science Centre  
All data and info are at [https://stereo-ssc.nascom.nasa.gov/beacon/beacon\\_secchi.shtml](https://stereo-ssc.nascom.nasa.gov/beacon/beacon_secchi.shtml)

Twin spacecraft in sun orbit, one ahead (ST-A) of Earth in its orbit, the other trailing behind (ST-B). They were launched on 25 October 2006.

Contact with ST-B was lost on 1 October 2014 during a test. Re-establishing contact will be tried again in June 2017.

Both spacecraft have passed the anti-solar point in 2015, heading back to earth, but now ST-A is trailing the earth (not re-labelled).

Spacecraft will be close to earth again somewhere in 2023.


The main SWx advantages of the STEREO mission are

- A stereoscopic view of the Sun and CMEs, allowing e.g. a much better determination of the speed and direction of a CME.
- A direct view on the Sun's farside and its solar activity.
- Knowing 1-2 weeks in advance on the strength of CH HSS or active sunspot groups that may rotate over the east limb as seen from earth.

There are 4 main instrument packages:

- SECCHI: Sun Earth Connection Coronal and Heliospheric Investigation
  - EUVI: extreme ultraviolet imager (4 passbands: 171, 195, 284, 304 Angstrom or 17.1, 19.5, 28.4, 30.4 nm)
  - COR1/2: white-light coronagraphs (inner: 1.3 to 4 solar radii/outer: 2-15 solar radii)
  - HI1/2: heliospheric imagers
- PLASTIC: PLASma and SupraThermal Ion Composition
  - Specifically interesting for proton and SEP events
- IMPACT: In-situ Measurements of Particles and CME Transients
  - Solar wind measurements
- SWAVES: STEREO Waves instrument
  - Track radiobursts from the Sun

# STEREO




The example is from the Carrington-like event from 23 July 2012 (Sun's farside), showing:  
Previous slide

- The EUVI304, COR1 and COR2 shortly after the maximum of the flare (but before the start of the proton event);
- The Type II and III radiobursts associated to the event (SWAVES)

This slide:

- The solar wind data from from PLASTIC/IMPACT for 7 days centered on 23 July 2012. Notice the strength of the event as observed by ST-A, and the lack of impact as observed by ST-B which was at the other side of the Sun
- The SEP data from the IMPACT instrument for 7 days centered on 23 July 2012. Notice the obvious differences between a well-connected (ST-A) and a poorly connected (ST-B) observer (much later, gradual and weaker with ST-B).

## STEREO orbit (2006-2022)



SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



[https://stereo-ssc.nascom.nasa.gov/cgi-bin/make\\_where\\_gif](https://stereo-ssc.nascom.nasa.gov/cgi-bin/make_where_gif)

## Exercise: Space-based instruments

- You want to have a clear view of any coronal holes that are visible on the Sun. Which instrument(s) would you use?
  - SOHO / LASCO C2
  - STEREO / Waves
  - SDO / AIA 193
  - GOES / XRS



:Issued: 2014 Apr 17 1325 UTC  
:Product: documentation at <http://www.sidc.be/products/tot>  
#-----#  
# DAILY BULLETIN ON SOLAR AND GEOMAGNETIC ACTIVITY from the SIDC #  
#-----#  
SIDC URSIGRAM 40417  
SIDC SOLAR BULLETIN 17 Apr 2014, 1304UT

SIDC FORECAST (valid from 1230UT, 17 Apr 2014 until 19 Apr 2014)  
SOLAR FLARES : Active (M-class flares expected, probability >=50%)  
GEOMAGNETISM : Quiet (A<20 and K<4)  
SOLAR PROTONS : Quiet

PREDICTIONS FOR 17 Apr 2014 10CM FLUX: 180 / AP: 013  
PREDICTIONS FOR 18 Apr 2014 10CM FLUX: 184 / AP: 007  
PREDICTIONS FOR 19 Apr 2014 10CM FLUX: 188 / AP: 005

COMMENT: Eleven sunspot groups were reported by NOAA today. NOAA ARs 2035, 2036, and 2037 (Catania numbers 24, 25, and 26 respectively) maintain the beta-gamma configuration of the photospheric magnetic field. The strongest flare of the past 24 hours was the M1.0 flare peaking at 19:59 UT yesterday in the NOAA AR 2035 (Catania number 24). The flare was associated with an EIT wave and a weak coronal dimming, but the associated CME was narrow and is not expected to arrive at the Earth.

We expect further flaring activity on the C-level, especially in the NOAA ARs 2035 and 2037 (Catania numbers 24 and 26 respectively) as well as in the NOAA AR 2042 (no Catania number yet) that yesterday appeared from behind the east solar limb, with a good chance for an M-class event.

Since yesterday evening the Earth is situated inside a solar wind structure with an elevated interplanetary magnetic field magnitude (occasionally up to 10 nT). It may be a weak ICME or the compression region on the flank of an ICME that missed the Earth. The solar origin of this structure is not clear. The north-south magnetic field component Bz was not strong, so no significant geomagnetic disturbance resulted (K index stayed below 4). Currently the solar wind speed is around 380 km/s and the IMF magnitude is around 8 nT.

We expect quiet to unsettled (K index up to 3) geomagnetic conditions, with active geomagnetic conditions (K = 4) possible, but unlikely.

TODAY'S ESTIMATED ISN : 145, BASED ON 17 STATIONS.  
99999

SOLAR INDICES FOR 16 Apr 2014  
WOLF NUMBER CATANIA : ///  
10CM SOLAR FLUX : 184  
AK CHAMBON LA FORET : 012  
AK WINGST : 004  
ESTIMATED AP : 004  
ESTIMATED ISN : 139, BASED ON 29 STATIONS.

NOTICEABLE EVENTS SUMMARY  
DAY BEGIN MAX END LOC XRAY OP 10CM Catania/NOAA RADIO\_BURST\_TYPES  
16 1954 1959 2004 S14E09 M1.0 1N 24/2035 II/2  
END



*Finding your way  
in the  
URSIGram*

*Satellites and instruments*

# Contents


- **Groundbased sensors**

- Visible light
- Radio domain
  - Humain
- Magnetosphere-Ionosphere
- Geomagnetism
- Neutron monitors
  - Dourbes

- **Spacebased sensors**

- GOES
- SDO
- PROBA2
- SOHO
- ACE
- DSCOVR
- STEREO

- Tools
- Overviews



### Space weather tools

Jan Janssens

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI

56


ESA Jhelioviewer: [http://swhv.oma.be/user\\_manual/](http://swhv.oma.be/user_manual/)



# Tools

- Various

- Soteria
- Solar Demon
- CACTus
- Drag model
- JHV (SWHV)
- STAFF
- COR2 J-plots
- COMESEP
- *10.7cm radioflux*
- *EUHFORIA*



**Soteria:** <http://www.sidc.be/soteria/soteria.php>

**Solar Demon:** <http://solardemon.oma.be/flares.php>

**CACTus:** <http://www.sidc.oma.be/cactus/out/latestCMEs.html>

**Drag model:** <http://oh.geof.unizg.hr/DBM/dbm.php>

**JHV (SWHV):** <http://www.jhelioviewer.org/>

**STAFF:** <http://www.staff.oma.be/>

**COR2 J-plots:** [http://www.sidc.be/rwc/cor2speed/cor2speed.html#canvas\\_position](http://www.sidc.be/rwc/cor2speed/cor2speed.html#canvas_position)

**COMESEP:** <http://www.comesep.eu/alert/>

10.7cm radioflux: will not be developed due to military interference

### EUFORIA: operational testing/development phase

**Other:**

Solar Monitor: <https://www.solarmonitor.org/index.php>

SWPC synoptic diagram: <https://www.swpc.noaa.gov/products/solar-synoptic-map>

Spacecast: <http://fp7-spacecast.eu/>

EURISGIC: <http://eurisgic.org/>

# Overviews

Forecast Weekly Presto Cactus All quiet CME arrival Monthly bulletin Quarterly Links

UTC time: 16:07:39

[General](#) | [Regions](#) | [Flares](#) | [Energetic Particles](#) | [Radio](#) | [CME](#) | [Solar Wind](#) | [Geomagnetism](#) | [Forecast Centres](#) | [Custom Campaigns](#)

## General

- [STAFF viewer \(backup\)](#)
- [Helioviewer \(backup\)](#)
- [Internal Documents](#) (webserver version)
- [cloud-as](#) (access to internal documents should be under SWOP\_InternalOperations)
- [password storage server](#) (in case forecaster needs passwords to IT infrastructure)

[Back to top](#)

## Regions

- [Soteria](#)
- [Solar Monitor Regions](#)
- [Latest NOAA synoptic map](#)
- [Rahen Maps](#)
- [STEREO Stormhurst heliographic maps](#)
- [PROBA2 LYRA data and SWAP images](#)
- [SDO movies](#)
- [SDO movies \(ROB page\)](#)
- [SOHO movies](#)
- [STEREO movies](#)
- [Realtime GONG H-alpha movies \(backup\)](#)

[Back to top](#)



SWx Dashboards in the SIDC SWx Forecast room

SWIC – Collaboration between

<http://sidc.be/previweb/links>

58




# Overviews

*Space Weather Now!*

<a href="#">Latest Reports</a>	<a href="#">Proton, SEP &amp; Radiation</a>	<a href="#">Solar Wind</a>
<a href="#">X-ray flares</a>	<a href="#">Radio Observations</a>	<a href="#">Geomagnetic</a>
<a href="#">X-ray EUV imagery</a>	<a href="#">CME</a>	<a href="#">Aurora</a>
<a href="#">White Light</a>	<a href="#">Coronal Holes</a>	<a href="#">Radio Comms &amp; TEC</a>
<a href="#">Magnetograms</a>	<a href="#">Farside</a>	<a href="#">Electrons</a>
<a href="#">H-alpha &amp; Ca II K</a>	<a href="#">Tools &amp; Data</a>	<a href="#">GIC</a>
<a href="#">Movie Centre</a>	<a href="#">Products</a>	<a href="#">Satellites</a>

## Current Space Weather Conditions




SWIC – Col

<http://users.telenet.be/j.janssens/Spaceweather.html>

59



# Overviews



<https://www.swpc.noaa.gov/>

SWIC – Collaboration between STCE, Koninklijke Luchtmacht, KNMI



NOAA/SWPC has developed SWx dashboards for specific SWx user communities such as aviation or satellites.

<https://www.swpc.noaa.gov/>

# Overviews

[illegible]

Other websites:

WMO/ICTSW: <https://www.wmo-sat.info/product-access-guide/theme/space-weather>

# Summary

- Both ground- and space-based data and imagery are used in SWx
  - Depends on the SWx user community and the purpose
- Multiple stations are a must
  - Back-up, cross-check & continuous monitoring
- Tools for analysis and forecasting are available
- Dashboards are available online